

Children's **TYLENOL**[®]

- The #1 branded fever reducer recommended by Pediatricians¹
- The fever reducer medicine with 0-6 month professional dosing



| Dose | | mL | Teaspoon (tsp)* | Tablet | Tablet |
|--------------------------|-----------|---------------------------------|------------------------------------|--------|--------|
| WEIGHT | AGE | (Use only the dropper provided) | (Use only the dosing cup provided) | | |
| 6-11 lbs | 0-3 mos | 0.4mL | | | |
| 12-17 lbs | 4-11 mos | 0.8mL | 1/2 (tsp) | | |
| 18-23 lbs | 12-23 mos | 1.2mL (0.8 + 0.4mL) | 3/4 (tsp) | | |
| 24-35 lbs | 2-3 yrs | 1.6mL (0.8 + 0.8mL) | 1 (tsp) | 2 | |
| 36-47 lbs | 4-5 yrs | | 1-1/2 (tsp) | 3 | |
| 48-59 lbs | 6-8 yrs | | 2 (tsp) | 4 | 2 |
| 60-71 lbs | 9-10 yrs | | 2 1/2 (tsp) | 5 | 2 1/2 |
| 72-95 lbs | 11 yrs | | 3 (tsp) | 6 | 3 |
| 96 lbs & over | 12 yrs | | | | 4 |

Use only as directed.

NOTE: If possible, use weight to dose; otherwise use age. To arrive at the correct dose, weigh your child before giving **TYLENOL**[®]. All doses may be repeated every 4 hours, but not more than 5 times daily.

A healthcare professional should be consulted for dosing for children under the age of two years.

WARNINGS:

- Children's **TYLENOL**[®] should not be taken for pain for more than 5 days or for fever for more than 3 days unless directed by a physician. If pain or fever persists or gets worse, if new symptoms occur, or if redness or swelling is present, a physician should be consulted because these could be signs of a serious condition.
- Do not exceed recommended dose. Taking more than the recommended dose (overdose) may not provide more relief and could cause serious health problems. Keep this and all drugs out of the reach of children. In case of accidental overdose, contact a physician or poison control center immediately. Prompt medical attention is critical even if you do not notice any signs or symptoms.
- Do not use with any other product containing acetaminophen.
- Do not use Adult Strength **TYLENOL**[®] products for children under 12 years of age.
- Do not use a kitchen teaspoon. Only use the dosing cup provided.

Children's **Motrin**[®]

- Pediatricians' #1 choice for fever relief that lasts up to 8 hours¹
- The #1 ibuprofen choice of consumers by more than 3 to 1 over Children's Advil[®] 2



| Dose | | mL | Teaspoon (tsp)* | Tablet | Tablet | Caplet |
|------------------------------------|-----------|---------------------------------|------------------------------------|-----------|---------------|---------------|
| WEIGHT | AGE | (Use only the dropper provided) | (Use only the dosing cup provided) | | | |
| Consult Your Child's Doctor | | | | | | |
| 12-17 lbs. | 6-11 mos | 1.25 mL | --- | --- | --- | --- |
| 18-23 lbs. | 12-23 mos | 1.875 mL | --- | --- | --- | --- |
| 24-35 lbs. | 2-3 yrs | --- | 1 (tsp) | 2 tablets | --- | --- |
| 36-47 lbs. | 4-5 yrs | --- | 1-1/2 (tsp) | 3 tablets | --- | --- |
| 48-59 lbs. | 6-8 yrs | --- | 2 (tsp) | 4 tablets | 2 tablets | 2 caplets |
| 60-71 lbs. | 9-10 yrs | --- | 2-1/2 (tsp) | 5 tablets | 2-1/2 tablets | 2-1/2 caplets |
| 72-95 lbs. | 11 yrs | --- | 3 (tsp) | 6 tablets | 3 tablets | 3 caplets |

One Dose Lasts 6-8 Hours

Children's Motrin is available over-the-counter (OTC) for the temporary reduction of fever and relief of minor aches and pains due to colds, flu, sore throat, headaches and toothaches. OTC Dosing: The recommended dose is 7.5 mg/kg every 6-8 hours. The recommended maximum daily dose is 30 mg/kg.

Keep all medications out of the reach of children. Use only as directed.

* Do not use a kitchen teaspoon. Only use the dosing cup provided.



McNEIL CONSUMER & SPECIALTY PHARMACEUTICALS
DIVISION OF McNEIL - PCC, INC.
FORT WASHINGTON, PA 19034 USA

References: 1. IMS NDI Pediatricians recommendations, 12 months ending August 2002
2. IRI Data, 52 weeks ending September 22, 2002.